








# Situation Analysis Dashboard

## National status Vs. two regions of Uganda

**Severity:**  
 Low  
 Medium  
 High

**Trend:**  
 Improvement  
 No change  
 Worsening

		Indicator	Status National	Trend	Severity	Target 2016	Status Western	Status Eastern
Nutritional Impact	Stunting	Prevalence of stunting among children <5 years old	33%			32%	44%	25%
	Wasting	Prevalence of wasting among children <5 years old	5%			N/A	3%	5%
	Underweight	Prevalence of underweight among children <5 years old	14%			10%	16%	10%
		Prevalence of underweight among non-pregnant women 15-49 years old (with BMI < 18.5 kg/m2)	12%			8%	8%	20%
	Iron deficiency	Prevalence of anaemia among children <5 years old	49%			50%	39%	55%
		Prevalence of anaemia among women 15-49 years old	23%			30%	17%	28%
Underlying Causes	Food Security	Percentage of households with poor or borderline food consumption	20%			N/A	18%	24%
	Health	Percentage of newborns weighing <2.5 kg at birth	10%			9%	8%	7%
	Care	Percentage of infants exclusively breastfed to age 6 months	63%			75%	???	???
		Prevalence of diarrhoea among children 6-59 months old	23%			N/A	19%	33%
Basic Causes	Education	Female literacy rate	64%			N/A	63%	49%
	Gender	Women's intra-household decision-making power	37%	N/A		N/A	37%	26%

Note: Statistics presented in red fall short of the established targets, whereas those presented in green exceed such targets.

Sources: DHS (2011 & 2006) / CFSVA (2013 & 2009)